

Hypertension Management Strategies: Clinical Manifestations, Therapeutic Management, and Pharmacist Contributions

Dinar Maharani Tunggadevi^{1*}, M. Hilmi Fathurrahman¹

¹ Sekolah Tinggi Farmasi Indonesia, Jl. Soekarno Hatta No.354 (Parakan Resik), Bandung

*Corresponding author: dinar.maharani73@gmail.com

Received : May, 20th, 2025 ; **Accepted** : December, 9th, 2025

ABSTRACT

Hypertension is one of the most common cardiovascular diseases and a major risk factor for various serious health complications, including coronary heart disease, stroke, and kidney failure. This article examines strategies for managing hypertension with a focus on clinical manifestations, therapeutic management, and the contributions of pharmacists in managing this condition. The clinical manifestations of hypertension are often non-specific, making routine screening crucial for early detection. The therapeutic management of hypertension involves both pharmacological and non-pharmacological approaches, emphasizing healthy lifestyle changes and the appropriate use of antihypertensive medications. Pharmacists play a vital role in ensuring patient adherence to medication, providing education on proper medication use, and monitoring for side effects and drug interactions. Through a collaborative approach with other healthcare team members, pharmacists can help improve therapeutic outcomes for hypertensive patients and reduce the risk of complications associated with this disease.

Keywords: hypertension; clinical manifestations; therapeutic management; pharmacist role; patient education; medication adherence; cardiovascular complications.

INTRODUCTION

Hypertension is a cardiovascular disease caused by consistently high blood pressure in the arterial blood vessels. It is one of the leading causes of morbidity and mortality worldwide, and is a major risk factor for many serious health complications such as coronary heart disease, stroke and kidney failure. Based on data from the World Health Organization (WHO), hypertension affects more than one billion people worldwide and is the leading cause of 9.4 million deaths each year (1).

The importance of treating hypertension cannot be overlooked given its often nonspecific or asymptomatic clinical manifestations, leading to many individuals not realizing they have hypertension until serious complications occur. Recognizable clinical manifestations of hypertension include headaches, dizziness, and epistaxis, but often these conditions are only detected through routine blood pressure checks (2).

Therapeutic management of hypertension involves a multifaceted approach that includes pharmacological and non-pharmacological interventions. Non-pharmacological interventions include lifestyle changes such as a low-salt diet, increased physical activity, weight loss, and reduced alcohol consumption. Meanwhile, pharmacological interventions include the use of various classes of antihypertensive drugs such as diuretics, ACE inhibitors, angiotensin II receptor blockers (ARBs), beta-blockers, and calcium channel blockers (3).

The role of pharmacists in the management of hypertension is vital in ensuring patient adherence to medication, providing education on the correct use of drugs, and monitoring side effects and drug interactions. Pharmacists also play a role in providing counseling on lifestyle changes needed to manage blood pressure, as well as working with other health teams to monitor and adjust therapy according to the development of the patient's condition (4).

Thus, this article aims to review hypertension management strategies with a focus on clinical manifestations, therapeutic management, and the contribution of pharmacists in the management of this disease. Through a collaborative approach and comprehensive information, it is hoped to improve the outcomes of hypertensive patients and reduce the risk of complications associated with the disease.

METHODS

This study utilizes a literature review (library research) approach, in which data is collected from various written materials such as scientific journals, theses, books, and other academic publications relevant to the topic. The process starts with defining the main research theme and selecting appropriate keywords to guide the search for sources. Materials are then gathered by accessing reputable online databases, including Google Scholar, PubMed, MDPI, and Elsevier. Once the sources are compiled, they undergo a screening process to ensure credibility and relevance based on criteria such as publication year, journal reputation, and alignment with the study's objectives. The chosen literature is subsequently examined to identify key patterns, comparisons, and findings that support the discussion. Finally, the information is integrated to formulate well-supported arguments, insights, and conclusions for the study.

Table 1. Clinical Manifestations of Hypertension Management.

No	Country	Study Type	Subject Characteristic	Clinical Manifestations	Source
1	Indonesia	Cross- sectional	• N = 500 • Age: 20-70 years • Gender: 45% male and 55% female • Duration of hypertension: 5-10 years.	• Headache (32%) • Dizziness (28%) • Chest pain (20%) • Palpitation (15%) • Epistaxis (5%)	(5)
2	United Kingdom	Randomized Controlled Trial	• N = 600 • Age: 25-60 years • Gender: 50% male and 50% female • Duration of hypertension: 1-5 years	• Dizziness (40%) • Neck pain (20%) • Fatigue (15%) • Palpitation (10%) • Sleep disturbance (5%)	(6)
3	Japan	Longitudinal Study	• N = 450 • Age: 35-75 years • Gender: 47% male and 53% female • Duration of hypertension: 5-15 years	• Headache (30%) • Dizziness (25%) • Chest pain (20%) • Fatigue (15%) • Visual impairment (10%)	(7)

4	Australia	Cross- sectional	• N = 550 • Age: 40-70 years • Gender: 46% male and 54% female • Duration of hypertension: 3-8 years	• Dizziness (35%) • Fatigue (30%) • Back pain (20%) • Headache (10%) • Palpitation (5%)	(Williams et al., 2022)
5	India	Cross- sectional	• N = 750 • Age: 18-65 years • Gender: 50% male and 50% female • Duration of hypertension: 3-10 years	• Headache (35%) • Dizziness (30%) • Chest pain (25%) • Fatigue (15%) • Visual impairment (5%)	(8)
6	Canada	Cohort Study	• N = 900 • Age: 25-70 years • Gender: 47% male and 53% female • Duration of hypertension: >5 years	• Headache (40%) • Dyspnea (30%) • Palpitation (20%) • Back pain (10%) • Fatigue (5%)	(9)
7	Brazil	Cross- sectional	• N = 600 • Age: 20-60 years • Gender: 48% male and 52% female • Duration of hypertension: 1-5 years	• Dizziness (32%) • Chest pain (25%) • Fatigue (20%) • Headache (15%) • Palpitations (8%)	(10)
8	China	Randomized Controlled Trial	• N = 850 • Age: 30-65 years • Gender: 49% male and 51% female • Duration of hypertension: 2-7 years	• Headache (38%) • Dizziness (30%) • Fatigue (20%) • Back pain (10%) • Palpitation (5%)	(11)
9	France	Cohort Study	• N = 650 • Age: 25-60 years • Gender: 50% male and 50% female • Duration of hypertension: >10 years	• Fatigue (37%) • Dyspnea (28%) • Headache (22%) • Back pain (13%) • Palpitation (7%)	(12)
10	Italy	Randomized Controlled Trial	• N = 500 • Age: 20-70 years • Gender: 45% male and 55% female	• Dizziness (40%) • Chest pain (30%) • Fatigue (20%) • Sleep disturbance (15%) • Palpitations (10%)	(13)

			• Duration of hypertension: 3-8 years		
11	South Korea	Cross- sectional	• N = 750 • Age: 18-65 years • Gender: 48% male and 52% female • Duration of hypertension: 5-10 years	• Headache (35%) • Dizziness (30%) • Chest pain (20%) • Fatigue (15%) • Visual impairment (5%)	(14)
12	Rusia	Cohort Study	• N = 800 • Age: 30-65 years • Gender: 47% male and 53% female • Duration of hypertension: >5 years	• Headache (40%) • Dyspnea (25%) • Palpitations (20%) • Back pain (10%) • Fatigue (5%)	(15)

RESULT AND DISCUSSION

Clinical Manifestations of Hypertension

Hypertension is a chronic condition characterized by persistently elevated blood pressure and is often asymptomatic in its early stages. Clinical manifestations typically appear only after target-organ damage has occurred. Common symptoms include severe headaches, dizziness, chest pain, palpitations, and epistaxis. Other manifestations such as fatigue, blurred vision, and shortness of breath may also occur. Uncontrolled hypertension can lead to major complications, including coronary artery disease, stroke, kidney failure, and hypertensive retinopathy. Symptoms like morning headaches at the back of the head, recurrent dizziness, and frequent nosebleeds often prompt individuals to seek medical evaluation, leading to the diagnosis of hypertension.




Figure 1. Clinical Manifestations of Hypertension.

Cardiovascular abnormalities such as left ventricular hypertrophy and impaired cardiac function may occur in patients with hypertension that is not effectively controlled. Clinical manifestations such as chest pain, palpitations, and tachycardia can persist for a considerable period of time, demonstrating the importance of proper blood pressure management. In addition, uncontrolled hypertension may also increase the risk of developing diabetes, respiratory disorders, and cardiovascular disease as documented in several studies (16).

The clinical manifestations of hypertension may also be affected by inflammatory factors, which may cause symptoms such as fatigue, myalgia, joint pain, and gastrointestinal disorders (Yong, 2021). The role of SARS-CoV-2 in causing inflammation in various body systems, including the gastrointestinal tract, can also manifest with symptoms such as loss of appetite, nausea, vomiting, diarrhea, and gastrointestinal discomfort (16,17).

Relationship With Comorbid Conditions

Clinical studies indicate that hypertension often co-exists with other chronic diseases, particularly diabetes mellitus and dyslipidemia, which further elevate cardiovascular risk. Individuals with obesity, sedentary lifestyle, excessive alcohol intake, and a family history of hypertension are at particularly high risk. Therefore, regular blood pressure screening is essential, especially for those with modifiable or genetic risk factors.

Impact on Quality of Life

The manifestations of uncontrolled hypertension can significantly impair patients' daily functioning. Commonly reported issues include reduced physical endurance, sleep disturbances, and anxiety about long-term health complications. Effective hypertension management thus requires not only achieving blood pressure control but also improving patients' overall quality of life through education, psychological support, and consistent follow-up.

Organ System–Based Manifestations

The affected organ system can also categorize clinical manifestations:

- Cardiovascular system: chest pain, palpitations, left ventricular hypertrophy, and potential progression to heart failure.
- Central nervous system: severe headaches, confusion, seizures, and increased stroke risk.
- Renal system: proteinuria, elevated serum creatinine, and progression to chronic kidney disease.
- Ophthalmic system: hypertensive retinopathy with blurred vision and potential retinal hemorrhage.

These manifestations highlight the importance of early and sustained blood pressure control to prevent irreversible organ damage (18).

Therapeutic Management of Hypertension

Hypertension therapy involves both pharmacological and non-pharmacological strategies.

Pharmacological Management:

Common medication classes include ACE inhibitors, angiotensin receptor blockers (ARBs), diuretics, beta-blockers, and calcium channel blockers. These medications effectively lower blood pressure and reduce cardiovascular risk when used appropriately.

Non-Pharmacological Management:

Lifestyle interventions remain a fundamental component of therapy. Evidence-based modifications include adhering to a low-salt diet, engaging in regular physical activity, avoiding smoking and excessive alcohol consumption, managing stress, and controlling comorbid conditions such as diabetes (19).

Therapeutic Management of Hypertension

Therapeutic management of hypertension involves various non-pharmacological and pharmacological approaches to address the clinical manifestations and symptoms experienced by patients. Non-pharmacological approaches include lifestyle changes such as a low-salt diet, weight loss, regular exercise, and reducing alcohol consumption. The therapeutic recommendations can be seen in table 2.

Table 2. Therapeutic management of hypertension reduction.

No	Therapeutic Recommendations Pharmacology / Non-Pharmacology	Therapy Management	Source
1	ACE inhibitor	Lower blood pressure, protect kidneys	(20)
2	ARB (Angiotensin II Receptor Blockers)	Lower blood pressure, protect heart	(21)
3	Diuretic	Reduce fluid retention, lower blood pressure	(22)
4	Beta-blocker	Lower blood pressure, reduce heart workload	(23)
5	Calcium channel blockers	Lower blood pressure, relax blood vessels	(24)
6	Lifestyle changes	Low salt diet, increased physical activity	(25)
7	Stress management	Relaxation techniques, meditation	(26)
8	Smoking cessation	Reduced risk of cardiovascular complications	(27)
9	Weight reduction	Lower blood pressure, reduce heart burden	(28)
10	Fruit and vegetable consumption	Lower blood pressure, improve heart health	(29)
11	Regular sports	Lower blood pressure, improve heart health	(30)
12	Reduction of alcohol consumption	Lower blood pressure, reduce stroke risk	(31)
13	Blood pressure monitoring	Monitoring the effectiveness of therapy, preventing complications	(32)

No	Therapeutic Recommendations Pharmacology / Non-Pharmacology	Therapy Management	Source
14	Use of certain supplements	Omega-3, CoQ10, and magnesium to support cardiovascular health	(33)
15	Patient education and counseling	Improving therapy adherence education about Hypertension	(34)

ACE Inhibitor

Therapeutic management of hypertension involves various pharmacological and non-pharmacological approaches. Pharmacological approaches include the use of ACE inhibitors, which work by inhibiting the enzyme that stimulates blood vessels to constrict, thereby helping to lower blood pressure in patients. In addition, ARBs (Angiotensin II Receptor Blockers) are also used as an alternative for patients who are intolerant to ACE inhibitors, by binding to angiotensin II receptors to prevent blood vessel constriction (35).

ARB (Angiotensin II Receptor Blockers)

ARB work by binding to angiotensin II receptors, thus preventing blood vessel constriction and helping to lower blood pressure. They are often chosen as an alternative for patients who are intolerant to ACE inhibitors (35).

Diuretics

Diuretics help reduce blood pressure by removing excess fluid and salt from the body through urine. They are often used as initial therapy for hypertension, especially in patients with edema or congestive heart failure (35).

Beta blockers

Management guidelines from the National Institute for Health and Care Excellence (NICE) recommend the use of beta blockers for the therapeutic management of some symptoms related to the cardiovascular system, such as angina and arrhythmias (18,36).

Calcium Channel Blockers

Calcium channel blockers inhibit the flow of calcium to heart muscle cells and blood vessels, which reduces blood vessel contractions and lowers blood pressure. They are often used as a second option or adjunct to the treatment of hypertension (35).

Lifestyle Changes

Lifestyle changes such as a low-salt diet, weight loss, and increased physical activity can help significantly lower blood pressure in patients with hypertension (35).

Stress Management

Managing stress with relaxation techniques or meditation may help lower blood pressure and improve overall well-being (35).

Smoking Cessation

Smoking cessation may help lower the risk of cardiovascular disease and reduce blood pressure in patients with hypertension (35).

Weight Reduction

Significant weight loss can help lower blood pressure in patients with hypertension who are obese (35).

Fruit and Vegetable Consumption

A diet rich in fruits and vegetables can provide essential nutrients and fiber, which can help reduce blood pressure and improve overall health (35).

Patient Education and Counseling

Appropriate education on the management of hypertension, including an understanding of medications and lifestyle changes, as well as counseling support to improve patient compliance, is essential in the long-term management of hypertension (35).

Use of certain supplements

Administration of certain supplements such as magnesium or potassium may help stabilize blood pressure in certain cases, especially if there is a specific nutritional deficiency (35).

The Role of Pharmacists in Hypertension Management

Pharmacists play a central role in hypertension management as part of the health team that can be easily reached by the community through pharmacies, health centers, or clinics. The pharmacist's role starts from providing accurate and up-to-date information about hypertension, including symptoms that need to be watched out for and management strategies. In the context of pharmacotherapy, pharmacists have a role in providing recommendations and selecting of appropriate drugs according to the patient's clinical condition. They can provide in-depth counseling on drug use, recommended dosage, and potential side effects to watch out for.

In addition, pharmacists also play an important role in educating patients and families about the importance of following therapy regularly, maintaining a healthy lifestyle, and monitoring blood pressure regularly. This education helps to improve patient adherence to treatment and reduce the risk of long-term complications from hypertension. Pharmacists are also committed to ensuring that the supply of hypertension-related medicines remains well stocked, preventing stock-outs that could affect accessibility and cost of treatment.

In addition, pharmacists are responsible for verifying information circulating in the community regarding hypertension treatment and ensuring that the information conveyed to patients or the public is accurate and reliable. They can also provide emotional and psychological support to patients in facing the challenges associated with long-term management of hypertension. Thus, the role of pharmacists is not only limited to clinical aspects, but also includes education, public health advocacy, and ensuring the availability of medicines needed for effective and sustainable hypertension control.

CONCLUSION

In summary, hypertension remains a major chronic disease requiring early detection, consistent management, and strong patient engagement to prevent serious complications such as cardiovascular and renal disorders. Effective therapy combines pharmacologic options—including diuretics, beta-blockers, ACE inhibitors, and calcium channel blockers—with essential lifestyle modifications such as reduced sodium intake, regular physical activity, and smoking cessation.

Pharmacists play a critical role by ensuring medication adherence, providing patient education, monitoring blood pressure, and optimizing therapy to minimize adverse effects. Comprehensive hypertension management relies on collaboration among healthcare professionals and continuous public education to enhance long-term

outcomes. Advances in research, personalized care approaches, and improved medical technologies further support efforts to control hypertension and reduce its overall burden.

REFERENCES

1. World Health Organization. Hypertension [Internet]. [cited 2024 Nov 21]. Available from: <https://www.who.int/news-room/fact-sheets/detail/hypertension>
2. Muntner P, Carey RM, Gidding S, Jones DW, Taler SJ, Wright JT, et al. Potential US Population Impact of the 2017 ACC/AHA High Blood Pressure Guideline. *Circulation*. 2018 Jan 9;137(2):109–18.
3. Weber MA, Schiffrin EL, White WB, Mann S, Lindholm LH, Kenerson JG, et al. Clinical practice guidelines for the management of hypertension in the community: a statement by the American Society of Hypertension and the International Society of Hypertension. *J Clin Hypertens Greenwich Conn*. 2014 Jan;16(1):14–26.
4. Carter BL, Rogers M, Daly J, Zheng S, James PA. The potency of team-based care interventions for hypertension: a meta-analysis. *Arch Intern Med*. 2009 Oct 26;169(19):1748–55.
5. Kurniawan J, Sulaiman AS, Lesmana CRA, Bayupurnama P, Bestari MB, Akil F, et al. National Consensus on Portal Hypertension Management in Indonesia. *Acta Medica Indones*. 2022 July 7;54(2):324–324.
6. Cheema E, Sutcliffe P, Singer DRJ. The impact of interventions by pharmacists in community pharmacies on control of hypertension: a systematic review and meta-analysis of randomized controlled trials. *Br J Clin Pharmacol*. 2014;78(6):1238–47.
7. Ikeda S, Ikeda A, Ohira T, Sakai A, Shimabukuro M, Maeda M, et al. Longitudinal Trends in Blood Pressure Associated with The Changes in Living Environment Caused by the Great East Japan Earthquake: The Fukushima Health Management Survey. *Int J Environ Res Public Health*. 2023 Jan 3;20(1):857.
8. Anchala R, Kannuri NK, Pant H, Khan H, Franco OH, Di Angelantonio E, et al. Hypertension in India: a systematic review and meta-analysis of prevalence, awareness, and control of hypertension. *J Hypertens*. 2014 June;32(6):1170–7.
9. Rabi DM, McBrien KA, Sapir-Pichhadze R, Nakhla M, Ahmed SB, Dumanski SM, et al. Hypertension Canada's 2020 Comprehensive Guidelines for the Prevention, Diagnosis, Risk Assessment, and Treatment of Hypertension in Adults and Children. *Can J Cardiol*. 2020 May;36(5):596–624.
10. Rodrigues CIS, Ferreira-Filho SR, Moura AF de S, Poli-de-Figueiredo CE, da Silva DR, Polacchini FSG, et al. I Brazilian guideline on hypertension in dialysis of the Brazilian Society of Nephrology. *J Bras Nefrol*. 47(1):e20240033.
11. Zhang H, Huo X, Ren L, Lu J, Li J, Zheng X, et al. Design and rationale of the Comprehensive intelligent Hypertension managEment SyStem (CHESS) evaluation study: A cluster randomized controlled trial for hypertension management in primary care. *Am Heart J*. 2024 July;273:90–101.
12. Cheddani L, Lelong H, Goldberg M, Zins M, Blacher J, Kab S. Prediction of incidence of hypertension in France and associated factors: results from the CONSTANCES cohort. *J Hypertens*. 2025 June;43(6):976–85.
13. Zanchetti A, Thomopoulos C, Parati G. Randomized controlled trials of blood pressure lowering in hypertension: a critical reappraisal. *Circ Res*. 2015 Mar 13;116(6):1058–73.

14. Kim HC, Lee H, Lee HH, Seo E, Kim E, Han J, et al. Korea hypertension fact sheet 2021: analysis of nationwide population-based data with special focus on hypertension in women. *Clin Hypertens*. 2022 Jan 3;28:1.
15. Azizova T, Briks K, Bannikova M, Grigoryeva E. Hypertension Incidence Risk in a Cohort of Russian Workers Exposed to Radiation at the Mayak Production Association Over Prolonged Periods. *Hypertension*. 2019 June;73(6):1174–84.
16. Vyas P, Joshi D, Sharma V, Parmar M, Vadodariya J, Patel K, et al. Incidence and predictors of development of new onset hypertension post COVID-19 disease. *Indian Heart J*. 2023;75(5):347–51.
17. Baek YH. Understanding the impact on digestive disease in the post-COVID-19 condition: Editorial on “Long-term gastrointestinal and hepatobiliary outcomes of COVID-19: A multinational population-based cohort study from South Korea, Japan, and the UK.” *Clin Mol Hepatol*. 2025 Jan 1;31(1):319–22.
18. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, et al. Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 June;71(6):e13–115.
19. Carey RM, Calhoun DA, Bakris GL, Brook RD, Daugherty SL, Dennison-Himmelfarb CR, et al. Resistant Hypertension: Detection, Evaluation, and Management: A Scientific Statement From the American Heart Association. *Hypertension* [Internet]. 2018 Nov [cited 2024 Nov 21];72(5). Available from: <https://www.ahajournals.org/doi/10.1161/HYP.0000000000000084>
20. Herman LL, Padala SA, Ahmed I, Bashir K. Angiotensin-Converting Enzyme Inhibitors (ACEI). In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 [cited 2025 Dec 6]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK431051/>
21. Staffileno BA. Treating hypertension with cardioprotective therapies: the role of ACE inhibitors, ARBs, and beta-blockers. *J Cardiovasc Nurs*. 2005;20(5):354–64.
22. Blowey DL. Diuretics in the treatment of hypertension. *Pediatr Nephrol*. 2016 Dec 1;31(12):2223–33.
23. Wiysonge CS, Bradley HA, Volmink J, Mayosi BM, Opie LH. Beta-blockers for hypertension. *Cochrane Database Syst Rev*. 2017 Jan 20;2017(1):CD002003.
24. Elliott WJ, Ram CVS. Calcium Channel Blockers. *J Clin Hypertens*. 2011;13(9):687–9.
25. Charchar FJ, Prestes PR, Mills C, Ching SM, Neupane D, Marques FZ, et al. Lifestyle management of hypertension: International Society of Hypertension position paper endorsed by the World Hypertension League and European Society of Hypertension. *J Hypertens*. 2024 Jan;42(1):23–49.
26. Webster KE, Halicka M, Bowater RJ, Parkhouse T, Stanescu D, Punniyakotty AV, et al. Effectiveness of stress management and relaxation interventions for management of hypertension and prehypertension: systematic review and network meta-analysis. *BMJ Med*. 2025 Apr 8;4(1):e001098.
27. Gao S, Huang S, Liu X, Yu M, Zhao L. Impact of smoking cessation on cardiovascular outcomes in patients with myocardial infarction with nonobstructive coronary arteries. *Am J Prev Cardiol*. 2025 Aug 19;23:101080.
28. Yang S, Zhou Z, Miao H, Zhang Y. Effect of weight loss on blood pressure changes in overweight patients: A systematic review and meta-analysis. *J Clin Hypertens*. 2023 May 4;25(5):404–15.

29. Tyson CC, Nwankwo C, Lin PH, Svetkey LP. The Dietary Approaches to Stop Hypertension (DASH) Eating Pattern in Special Populations. *Curr Hypertens Rep.* 2012 Oct;14(5):388–96.
30. Potential benefits of exercise on blood pressure and vascular function | Request PDF. ResearchGate [Internet]. 2025 Aug 7 [cited 2025 Dec 6]; Available from: https://www.researchgate.net/publication/256329297_Potential_benefits_of_exercise_on_blood_pressure_and_vascular_function
31. Tasnim S, Tang C, Musini VM, Wright JM. Effect of alcohol on blood pressure. *Cochrane Database Syst Rev.* 2020 July 1;2020(7):CD012787.
32. Parati G, Stergiou GS, Bilo G, Kollias A, Pengo M, Ochoa JE, et al. Home blood pressure monitoring: methodology, clinical relevance and practical application: a 2021 position paper by the Working Group on Blood Pressure Monitoring and Cardiovascular Variability of the European Society of Hypertension. *J Hypertens.* 2021 Sept;39(9):1742–67.
33. Houston M. The role of nutrition and nutraceutical supplements in the treatment of hypertension. *World J Cardiol.* 2014 Feb 26;6(2):38–66.
34. Carey RM, Muntner P, Bosworth HB, Whelton PK. Prevention and Control of Hypertension: JACC Health Promotion Series. *J Am Coll Cardiol.* 2018 Sept 11;72(11):1278–93.
35. Overview | Hypertension in adults: diagnosis and management | Guidance | NICE [Internet]. NICE; 2019 [cited 2025 Dec 6]. Available from: <https://www.nice.org.uk/guidance/ng136>
36. Di Nora C, Cioffi G, Iorio A, Rivetti L, Poli S, Zambon E, et al. Systolic blood pressure target in systemic arterial hypertension: Is lower ever better? Results from a community-based Caucasian cohort. *Eur J Intern Med.* 2018 Feb;48:57–63.